VVARDIS
SWITZERLAND

BIOMIMETIC DENTAL SCIENCE

CURODONT™ REPAIR FLUORIDE PLUS

Anticavity Therapy via Guided Enamel Remineralization

SWISS MADE

INCREASE PRODUCTION WITH CURODONT™ REPAIR FLUORIDE PLUS

Directions for Use – Workflow

VVARDIS
SWITZERLAND

BIOMIMETIC DENTAL SCIENCE

How to apply Curodont™ Repair Fluoride Plus

1. Perform oral prophylaxis, especially in the area to be treated to eliminate any calculus/ plaque/ materia alba.
2. Use cheek retractors and cotton rolls in the buccal and lingual vestibules adjacent to the lesion. Use of a rubber dam is optional.
3. Dip a small cotton pellet, in 2% sodium hypochlorite.
4. Use the saturated cotton pellet for 20 seconds to clean the lesion and remove the organic pellicle. (Figure 2)
5. Rinse with water.
6. Apply 35% phosphoric acid to the area in question for 20 seconds and then, rinse. Unwaxed dental floss may be used to distribute it in the interdental space. (Figure 3)
7. Gently dry the area to be treated.
8. Take one CURODONT™ REPAIR FLUORIDE PLUS applicator tip and press the sponge into the material. Withdraw the sponge and press it onto the lesion. For occlusal lesions, pressing the sponge is enough to distribute the material into the lesion; spreading it over the tooth is unnecessary. Use one applicator per lesion.
   - N.B. for adjacent interproximal lesions, one applicator will suffice. (Figure 4)
9. Keep isolation for 5 minutes. After which, the patient can spit if needed, but not rinse.
10. Discharge the patient, instructing not to rinse, eat or drink for 30 minutes and maintain good oral hygiene.

Buccal/Lingual Caries **Occlusal Caries** **Interproximal Caries**

Figure 1

B: Incipient enamel lesion on the labial surface of the central incisor.
O: Incipient enamel lesion on the occlusal surface of a mandibular first molar.
I: Incipient enamel lesion on the proximal surfaces of a mandibular first and second molar.

Figure 2



Apply 2% sodium hypochlorite for 20 seconds to remove the organic pellicle.
Rinse with water.

Figure 3



Apply 35% orthophosphoric acid for 20 seconds.

Figure 4

Apply CURODONT™ REPAIR FLUORIDE PLUS with the applicator tip, then wait 5 minutes.