

# vVARDIS

SWITZERLAND

BIOMIMETIC DENTAL SCIENCE



## CURODONT™ PROTECT Anti-cavity & Anti-sensitivity Dental Gel

### HOW TO APPLY CURODONT™ PROTECT IN-OFFICE

CURODONT™ PROTECT is quick and easy to apply with the help of a rubber polisher, interdental brush, or a micro-brush.

Before professional cleaning/bleaching or any invasive treatment

Apply 2 minutes before the dental procedure to reduce sensitivity and discomfort during the visit.

After treatment with  
CURODONT™ REPAIR FLUORIDE PLUS

Wait for 5 minutes and apply on treated areas and on the rest of the teeth.

### HOW TO APPLY CURODONT™ PROTECT AT HOME

CURODONT™ PROTECT can be applied with a toothbrush, interdental brush, or a clean finger.

After treatment with  
CURODONT™ REPAIR FLUORIDE PLUS

Once to twice a week, apply a thin layer over all teeth surfaces after brushing at bedtime.

For caries prevention

Once to twice a week, apply a thin layer over all teeth surfaces after brushing at bedtime.


For sensitivity relief

Once or several times a day until the pain subsides.

**Please note:** The gel must be allowed to stay on the teeth undisturbed for 1-2 minutes before spitting out the residue, if any. This should not be followed by rinsing to maintain the contact of the gel with the teeth for as long as possible.

### MAXIMIZE OUTCOMES WITH THE vVARDIS CURODONT™ SYSTEM FOR WATCH AREAS MANAGEMENT

**Treat** with  
CURODONT™ REPAIR FLUORIDE PLUS  
in-office



**Maintain & Protect** with  
CURODONT™ PROTECT  
in-office and/or at-home